


Application No. 10/511534
Reply to Office Action of March 3, 2006

AMENDMENTS TO THE CLAIMS


1. (Currently amended) Reactive dyes as per the hereinbelow indicated and defined general formula (I),



(I)

where

D¹ and D² are independently a group of the general formula (1)



(1)

where

R¹ and R² are independently hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl, sulfo, carboxyl, cyano, nitro, amido, ureido or halogen; and

X¹ is hydrogen or a group of the formula -SO₂-Z,

where


Z is -CH=CH₂, -CH₂CH₂Z¹ or hydroxyl,

where

Application No. 10/511534
Reply to Office Action of March 3, 2006

Z^1 is hydroxyl or an alkali-detachable group; or

D^1 and D^2 are independently a naphthyl group of the general formula (2)




where

R^3 and R^4 are independently hydrogen, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, hydroxyl, sulfo, carboxyl, cyano, nitro, amido, ureido or halogen; and

X^2 has one of the meanings of X^1 ; or

D^1 and D^2 are independently a group of the general formula (3)



where


R^5 and R^6 independently have one of the meanings of R^1 and R^2 ;

R^7 is hydrogen, (C_1-C_4) -alkyl, unsubstituted or (C_1-C_4) -alkyl-, (C_1-C_4) -alkoxy-, sulfo-, halogen- or carboxyl-substituted phenyl; and

~~Z^2 is a heterocyclic reactive radical; or~~

Application No. 10/511534
Reply to Office Action of March 3, 2006

Z² a group of the general formula (4) or (5) or (6)



where


V is fluorine or chlorine;

U¹, U² are independently fluorine, chlorine or hydrogen; and

Q¹, Q² are independently chlorine, fluorine, cyanamido, hydroxyl, (C₁-C₆)-alkoxy,

phenoxy, sulfophenoxy, mercapto, (C₁-C₆)-alkylmercapto, pyridino, carboxypyridino,

carbamoylpyridino or a group of the general formula (7) or (8)



where

R⁸ is hydrogen or (C₁-C₆)-alkyl, sulfo-(C₁-C₆)-alkyl or phenyl unsubstituted or

substituted by (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, sulfo, halogen, carboxyl, acetamido,


ureido;

R⁹ and R¹⁰ independently have one of the meanings of R⁸ or combine to form a cyclic

ring system of the formula -(CH₂)_j-, where j is 4 or 5, or alternatively

Application No. 10/511534
Reply to Office Action of March 3, 2006

-(CH₂)₂-E-(CH₂)₂-, where E is oxygen, sulfur, sulfo, -NR¹¹ where R¹¹ = (C₁-C₆)-alkyl;
W is phenylene which is unsubstituted or substituted by 1 or 2 substituents, such as
(C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, carboxyl, sulfo, chlorine, bromine, or is
(C₁-C₄)-alkylene-arylene or (C₂-C₆)-alkylene, which can be interrupted by oxygen,
sulfur, sulfo, amino, carbonyl, carboxamido, or is phenylene-CONH-phenylene, which is
unsubstituted or substituted by (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl, sulfo, carboxyl,
amido, ureido or halogen, or is naphthylene which is unsubstituted or substituted by 1 or
2 sulfo groups; and
Z is as defined above; or
 D¹ and D² are independently a group of the general formula (9)



where


R¹² is hydrogen, (C₁-C₄)-alkyl, aryl or a substituted aryl radical;

R¹³ and R¹⁴ are independently hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl,

sulfo, carboxyl, cyano, nitro, amido, ureido or halogen; and


A is a phenylene group of the general formula (10)

Application No. 10/511534
Reply to Office Action of March 3, 2006




where

R^{15} and R^{16} are independently hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl, sulfo, carboxyl, cyano, nitro, amido, ureido or halogen; or a naphthylene group of the general formula (11)



where

R^{17} and R^{18} are independently hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl, sulfo, carboxyl, cyano, nitro, amido, ureido or halogen; or a polymethylene group of the general formula (12)



where

k is a whole number greater than 1 and

R^{19} and R^{20} are independently hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, hydroxyl,

cyano, amido, halogen or aryl; and

X^3 has one of the meanings of X^1 ; and

Application No. 10/511534
Reply to Office Action of March 3, 2006

R, R* are independently hydrogen, (C₁-C₄)-alkyl or sulfomethyl; and

M is hydrogen, an alkali metal or one equivalent of an alkaline earth metal,
with the proviso that the dyes of the general formulae (I) contain at least one fiber-
reactive heterocyclic group of the general ~~formula~~ formula (4) or (6).

2. (Canceled)

3. (Previously presented) Reactive dyes as per claim 1, where the substituents R are
hydrogen or sulfomethyl and R* is hydrogen.

4. (Currently amended) Reactive dyes as per claim 1, ~~characterized in that~~ wherein the
substituents R¹ and R² are hydrogen, methyl, methoxy or sulfo, R¹² to R¹⁶ are hydrogen
and R³ to R⁶, R¹⁷ and R¹⁸ are hydrogen or sulfo, R⁷ and R⁸ are hydrogen methyl or
phenyl and R⁹ and R¹⁰ are hydrogen, methyl 2-sulfoethyl, 2-, 3- or 4-sulfophenyl, or R⁹
and R¹⁰ combine to form a cyclic ring system which conforms to the formula -(CH₂)₂-
O-(CH₂)₂-.


5. (Currently amended) Reactive dyes as per claim 1, ~~characterized in that~~ wherein Z is
vinyl, β-chloroethyl or β-sulfatoethyl.

6. (Currently amended) Reactive dyes as claimed in ~~claim 2~~ claim 1, ~~characterized in that~~
wherein Q¹ and Q² in the general formula (5) are independently fluorine, chlorine,
cyanamido, morpholino, 2-sulfophenylamino, 3-sulfophenylamino, 4-sulfophenylamino,


Application No. 10/511534
Reply to Office Action of March 3, 2006

3-(2-sulfatoethylsulfonyl)phenylamino, 4-(2-sulfatoethylsulfonyl)phenylamino,
3-(vinylsulfonyl)phenylamino, 4-(vinylsulfonyl)phenylamino, N-methyl-
N-(2-(2-sulfatoethylsulfonyl)ethyl)amino or N-phenyl-N-(2-(2-sulfatoethyl-
sulfonyl)ethyl)amino.

7. (Previously presented) A process for preparing dyes of the general formula (I) as per claim 1, which comprises diazotizing one equivalent of an amine of the general formula (16)




where D^1 is as defined in claim 1 and the resulting diazonium compound being reacted in a first stage with an aqueous solution or suspension of one equivalent of a coupling component as per the general formula (17)




where R, R* and M are each as defined in claim 1, to form a monoazo dye as per the general formula (13)

Application No. 10/511534
Reply to Office Action of March 3, 2006




(13)

and subsequently one equivalent of an amine of the general formula (18)




where D^2 is as defined in claim 1, being diazotized and the resulting diazonium compound being coupled in the second stage with the monoazo dye of the general formula (13) obtained in the first stage to form the disazo dye of the general formula (I).

8. (Currently amended) The process for preparing dyes of the general formula (I) as per claim 1 in the event that the groups D^1 and D^2 as per the general formulae (I) have the same meaning by two equivalents of an amine of the general formula (16)




where D^1 is as defined in claim 1 which comprises diazotizing and reacting in a first stage with one equivalent of a coupling component of the general formula (17)

Application No. 10/511534
Reply to Office Action of March 3, 2006



(17)

to form a monoazo dye of the general formula (13)



(13)

and the second coupling to the disazo dye of the general formula (I) where the groups D^1 and D^2 have the same meaning being carried out subsequently.

9. (Previously presented) An aqueous liquid preparation containing a dye as set forth in claim 1 at a level of 5-90% by weight.
10. (Canceled)
11. (Currently amended) Reactive dyes as per ~~claim 2~~ claim 1, where the substituents R are hydrogen or sulfomethyl and R^* is hydrogen.
12. (Currently amended) Reactive dyes as per claim 11, ~~characterized in that~~ wherein the substituents R^1 and R^2 are hydrogen, methyl, methoxy or sulfo, R^{12} to R^{16} are hydrogen

Application No. 10/511534
 Reply to Office Action of March 3, 2006

and R^3 to R^6 , R^{17} and R^{18} are hydrogen or sulfo, R^7 and R^8 are hydrogen methyl or phenyl and R^9 and R^{10} are hydrogen, methyl 2-sulfoethyl, 2-, 3- or 4-sulfophenyl, or R^9 and R^{10} combine to form a cyclic ring system which conforms to the formula $-(CH_2)_2-O-(CH_2)_2-$.

13. (Currently amended) Reactive dyes as per claim 12, ~~characterized in that~~ wherein Z is vinyl, β -chloroethyl or β -sulfatoethyl.
14. (Currently amended) Reactive dyes as claimed in claim 13, ~~characterized in that~~ wherein Q^1 and Q^2 in the general formula (5) are independently fluorine, chlorine, cyanamido, morpholino, 2-sulfophenylamino, 3-sulfophenylamino, 4-sulfophenylamino, 3-(2-sulfatoethylsulfonyl)phenylamino, 4-(2-sulfatoethylsulfonyl)phenylamino, 3-(vinylsulfonyl)phenylamino, 4-(vinylsulfonyl)phenylamino, N-methyl-N-(2-(2-sulfatoethylsulfonyl)ethyl)amino or N-phenyl-N-(2-(2-sulfatoethylsulfonyl)ethyl)amino.
15. (Previously presented) A process of dyeing or printing hydroxyl- and/or carboxamido-containing fiber material which comprises contacting the reactive dyes as per claim 1 with said material.